

Hemispheric-scale planktonic diatom (*Cyclotella*) response to recent warming

Kathleen Rühland*, Andrew M. Paterson#, and John P. Smol*


* (PEARL), Dept. of Biology, Queen's University, Kingston ON, CANADA, # Ontario Ministry of the Environment, Dorset ON, CANADA

Abstract


- A synthesis of over 200 diatom-based paleolimnological records from nonacidified/nonenriched lakes show remarkably similar climate-induced taxon-specific shifts since the 19th century across the Northern Hemisphere.
- These diatom shifts occurred in conjunction with changes in freshwater habitat structure and quality, which, in turn, we link to hemispheric warming trends.
- A marked increase in the relative abundances of planktonic *Cyclotella* taxa with concurrent sharp declines in thickly-silicified *Aulacoseira* taxa and/or benthic *Fragilaria* taxa were observed.
- Unparalleled warming over the last few decades resulted in substantial increases in the length of the ice-free period that, similar to 19th century changes in high-latitude lakes, likely triggered a reorganization of diatom community composition in lakes throughout North America and western Europe.
- Our case studies [Whitefish Bay (Lake of the Woods), Emmett & George lakes (Bruce Peninsula) Ontario] revealed that recent increases in the relative abundances of planktonic *Cyclotella* species and concurrent decreases in heavily silicified *Aulacoseira* and/or benthic *Fragilaria* species were strongly correlated to recent increases in temperature and substantially longer ice-free periods.

Meta-analysis

Timing of *Cyclotella* increases





Distribution of *Cyclotella* shift in N. Hemisphere




Case Studies

Whitefish Bay: Lake of the Woods, Ontario, Canada



The Kenora temperature data are available from <http://www.ccma.bc.ec.gc.ca/hccd>. The Whitefish Bay ice-out data were provided by the Ministry of Natural Resources, Kenora, Ontario, Canada.

Emmett & George Lakes: Bruce Peninsula, Ontario, Canada



Summary & Conclusions

- Diatom data from over 200 lakes provide a spatially coherent picture that substantial warming over the last few decades have driven taxon-specific changes.
 - Strikingly similar diatom shifts are now evident across vast regions of the Northern Hemisphere representing a wide spectrum of lake ecosystems.
 - Non-acidified, nutrient-poor, freshwater ecosystems throughout the Northern Hemisphere have crossed important climatically-induced ecological thresholds that were initiated in the 19th century in Arctic and alpine regions, but typically only occurred in the mid-20th century in lakes from mid-latitude regions of North America and western Europe.
 - The nature and timing of the taxon-specific diatom changes recorded in our meta-analysis is consistent with climate-related changes in the physical properties in lakes including:
 - Length of the open-water season
 - Timing, duration and strength of thermal stratification
 - Timing and strength of spring freshet
 - Timing and duration of spring and autumn overturn
 - Changes in these lake-water properties likely provide favourable habitats for small, fast-growing planktonic *Cyclotella* species such as *C. stelligera* complexes and *C. comensis* complexes. These planktonic diatoms may be able to exploit longer ice-free periods and/or deeper, subsurface habitats where nutrient concentrations are somewhat elevated and where light properties become more stabilized as thermal stratification develops (e.g. Fahnenstiel & Glime, 1983).
 - The recurring and widespread trend of recent increases in the relative abundances of planktonic *Cyclotella* species and concurrent decreases in heavily silicified *Aulacoseira* and/or benthic *Fragilaria* taxa represented by our three case studies were strongly correlated to recent increases in temperature and substantially longer ice-free periods.
 - Synergistic effects between climate warming and other human-induced stressors also occur. However, based on the timing, magnitude, and nature of the diatom shifts from this study, we conclude that warming-induced change in lake-ice cover (and associated limnological changes) was the primary explanatory metric for hemispheric-scale increases in planktonic *Cyclotella* species over the last ca. 200 years.
- If the rate and magnitude of temperature increase continues, it is likely that new ecological thresholds will be crossed, many of which may be unexpected.

Further reading

Rühland, K., Paterson, A.M., and Smol J.P. (2008). Hemispheric-scale patterns of climate-related shifts in planktonic diatoms from North American and European lakes. *Glob. Change Biol.* 14, doi: 10.1111

Smol, J.P. et al. (2005). Climate-driven regime shifts in the biological communities of arctic lakes. *Proc. Nat. Acad. Sci.* 102: 4397-4402.

This poster can be viewed as a pdf at <http://biology.queensu.ca/low/Research%20Page.html>